



**County of San Benito
Health & Human Services Agency
Public Health Services
Child Health and Disability Prevention Program**

Child Health and Disability Prevention Program Online Vision Screening Training



Objectives

By the end of the training, participants will be able to:

- Understand the CHDP requirements for vision screening
- Know the basic anatomy of the eye and the pathway of vision
- Perform a vision acuity screening on a preschool child
- Identify the correct screening charts and when to use them
- Document visual acuity screening results
- Refer to an eye specialist if needed

CHDP Visual Acuity Screening Requirements

- Screen for visual problems at every well child visit
- Visual Acuity Screening should be done beginning at age 3
- Conduct the screen in a well-lit room, free of visual and auditory distractions
- The eye chart should be at the child's eye level
- Each eye should be screened separately
- Proper selection of age-appropriate optotypes and testing methods are important in obtaining accurate screening results
- Screening distance is 10 feet. This short distance will enhance interaction between the child and screener without decreasing accuracy of screening results

CHDP Visual Acuity Screening Recommendations for Providers

- Complete a CHDP approved training in vision screening
 - Certificate is good for 4 years
- Either Critical Line or Threshold Screening may be used for the vision screening

Bright Futures Periodicity

American Academy of Pediatrics
DEDICATED TO THE HEALTH OF ALL CHILDREN

Recommendations for Preventive Pediatric Health Care
Bright Futures/American Academy of Pediatrics



Bright futures recommends risk assessments to be performed prior to the age of 3, visual acuity screenings to start at 3 till 6 years and at 8, 10, 12 and 15 years of age; risk assessment to be performed on the other years with appropriate interventions to follow, if positive results are found.

Each child and family is unique; therefore, these Recommendations for Preventive Pediatric Health Care are designed for the care of children who are receiving competent parenting, have no manifestations of any important health problems, and are growing and developing in a satisfactory fashion. Developmental, psychosocial, and chronic disease issues for children and adolescents may require frequent counseling and treatment visits separate from preventive care visits. Additional visits also may become necessary if circumstances suggest variations from normal.

These recommendations represent a consensus by the American Academy of Pediatrics (AAP) and Bright Futures. The AAP continues to emphasize the great importance of continuity of care in comprehensive health supervision and the need to avoid fragmentation of care. Refer to the specific guidance by age as listed in the *Bright Futures Guidelines* (Hagan JF, Shaw JS, Duncan PM, eds. *Bright Futures: Guidelines for Health Supervision of Infants, Children, and Adolescents*. 4th ed. Elk Grove Village, IL: American Academy of Pediatrics; 2017).

The recommendations in this statement do not indicate an exclusive course of treatment or standard of medical care. Variations, taking into account individual circumstances, may be appropriate. Copyright © 2017 by the American Academy of Pediatrics, updated February 2017. No part of this statement may be reproduced in any form or by any means without prior written permission from the American Academy of Pediatrics except for one copy for personal use.

	AGE ¹	PRENATAL ²	NEWBORN ³	3-5 d ⁴	1 mo	2 mo	4 mo	6 mo	9 mo	12 mo	15 mo	18 mo	24 mo	30 mo	3 y	4 y	5 y	6 y	7 y	8 y	9 y	10 y	11 y	12 y	13 y	14 y	15 y	16 y	17 y	18 y	19 y	20 y	21 y
HISTORY		•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•
MEASUREMENTS																																	
Length/Height and Weight			•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•
Head Circumference			•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•
Weight for Length			•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•
Body Mass Index ⁵			•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•
Blood Pressure ⁶			•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•
SENSORY SCREENING																																	
Vision ⁷		•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•
Hearing ⁸		•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•
DEVELOPMENTAL/BEHAVIORAL HEALTH																																	
Developmental Screening ⁹																																	
Autism Spectrum Disorder Screening ¹⁰																																	
Developmental Surveillance			•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•
Psychosocial/Behavioral Assessment ¹¹			•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•
Tobacco, Alcohol, or Drug Use Assessment ¹²																																	
Depression Screening ¹³																																	
Maternal Depression Screening ¹⁴																																	
PHYSICAL EXAMINATION¹⁵			•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•
PROCEDURES¹⁶																																	
Newborn Blood		•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•
Newborn Bilirubin ¹⁷		•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•
Critical Congenital Heart Defect ¹⁸		•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•
Immunization ¹⁹		•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•
Anemia ²⁰																																	
Lead ²¹																																	
Tuberculosis ²²																																	
Dyslipidemia ²³																																	
Sexually Transmitted Infections ²⁴																																	
HIV ²⁵																																	
Cervical Dysplasia ²⁶																																	
ORAL HEALTH²⁷																																	
Fluoride Varnish ²⁸																																	
Fluoride Supplementation ²⁹																																	
ANTICIPATORY GUIDANCE																																	

- If a child comes under care for the first time at any point on the schedule, or if any items are not accomplished at the suggested age, the schedule should be brought up-to-date at the earliest possible time.
- A prenatal visit is recommended for parents who are at high risk, for first-time parents, and for those who request a conference. The prenatal visit should include anticipatory guidance, pertinent medical history, and a discussion of benefits of breastfeeding and planned method of feeding, per "The Prenatal Visit" (<http://pediatrics.aappublications.org/content/124/4/227.full>).
- Newborns should have an evaluation after birth, and breastfeeding should be encouraged (and instruction and support should be offered).
- Newborns should have an evaluation within 3 to 5 days of birth and within 48 to 72 hours after discharge from the hospital to include evaluation for feeding and jaundice. Breastfeeding newborns should receive formal breastfeeding evaluation, and their mothers should receive encouragement and instruction, as recommended in "Breastfeeding and the Use of Human Milk" (<http://pediatrics.aappublications.org/content/120/3/487.full>). Newborns discharged less than 48 hours after delivery must be examined within 48 hours of discharge, per "Hospital Stay for Healthy Term Newborns" (<http://pediatrics.aappublications.org/content/125/2/405.full>).
- Screen, per "Expert Committee Recommendations Regarding the Prevention, Assessment, and Treatment of Child and Adolescent Overweight and Obesity: Summary Report" (http://pediatrics.aappublications.org/content/120/supplement_4/5164.full).

- Blood pressure measurement in infants and children with specific risk conditions should be performed at visits before age 3 years.
- A visual acuity screen is recommended at ages 4 and 5 years, as well as in cooperative 3-year-olds. Instrument-based screening may be used to assess risk at ages 12 and 24 months, in addition to the well visits at 3 through 5 years of age. See "Visual System Assessment in Infants, Children, and Young Adults by Pediatricians" (<http://pediatrics.aappublications.org/content/137/1/Ae20153596>) and "Procedures for the Evaluation of the Visual System by Pediatricians" (<http://pediatrics.aappublications.org/content/137/1/Ae20153597>).
- Confirm initial screen was completed, verify results, and follow-up, as appropriate. Newborns should be screened, per "Year 2007 Position Statement: Principles and Guidelines for Early Hearing Detection and Intervention Programs" (<http://pediatrics.aappublications.org/content/120/4/988.full>).
- Verify results as soon as possible, and follow-up, as appropriate.
- Screen with audiometry including 6000 and 8000 Hz high frequencies once between 11 and 14 years, once between 15 and 17 years, and once between 18 and 21 years. See "The Sensitivity of Adolescent Hearing Screens Significantly Improves by Adding High Frequencies" ([http://www.jahonline.org/article/S1054-139X\(16\)00048-3/fulltext](http://www.jahonline.org/article/S1054-139X(16)00048-3/fulltext)).
- See "Identifying Infants and Young Children With Developmental Disorders in the Medical Home: An Algorithm for Developmental Surveillance and Screening" (<http://pediatrics.aappublications.org/content/118/1/A05.full>).

- Screening should occur per "Identification and Evaluation of Children With Autism Spectrum Disorders" (<http://pediatrics.aappublications.org/content/120/5/1183.full>).
- This assessment should be family centered and may include an assessment of child social-emotional health, caregiver depression, and social determinants of health. See "Promoting Optimal Development: Screening for Behavioral and Emotional Problems" (<http://pediatrics.aappublications.org/content/135/2/289>) and "Poverty and Child Health in the United States" (<http://pediatrics.aappublications.org/content/137/4/Ae20160339>).
- A recommended assessment tool is available at <http://www.casat-boston.org/CRAFT/index.php>.
- Recommended screening using the Patient Health Questionnaire (PHQ-2) or other tools available in the GLAD-PC toolkit and at <http://www.aap.org/en-us/advocacy-and-policy/aap-health-initiatives/Mental-Health/Documents/Well-BeingChart.pdf>.
- Screening should occur per "Incorporating Recognition and Management of Perinatal and Postpartum Depression Into Pediatric Practice" (<http://pediatrics.aappublications.org/content/126/5/1032>).
- At each visit, age-appropriate physical examination is essential, with infant totally unclothed and older children undressed and suitably draped. See "Use of Chaperones During the Physical Examination of the Pediatric Patient" (<http://pediatrics.aappublications.org/content/127/5/991.full>).
- These may be modified, depending on entry point into schedule and individual need.

(continued)

KEY: • = to be performed ★ = risk assessment to be performed with appropriate action to follow, if positive ← • → = range during which a service may be provided

https://www.aap.org/en-us/Documents/periodicity_schedule.pdf

Please PAUSE and view
the videos located on
step 3